

PU-9536

Seat No.

Third Year B. Physiotherapy Examination July - 2018 General Surgery - 2

Time: 3 Hours] [Total Marks: 100

Instructions:

- 1. Write each section in separate answer book
- 2. Figures to the right indicate full marks
- 3. Write legibly and use illustrations where necessary

SECTION-1 Cardio Thoracic Surgery

1 Write answers: (Any Two)

20

- (a) Describe the causes pathophysiology, clinical features, investigations and treatment of Carcinoma of Lung.
- (b) Describe the causes pathophysiology, clinical features, investigations and treatment of Mitral Valve Stenosis (MS).
- (c) Describe the causes pathophysiology, clinical features, investigations and treatment of Pneumothorax.
- 2 Write short notes: (Any Two)

10

- (a) Cardiopulmonary bypass
- (b) Cervical rib
- (c) Felot's Tetrology
- 3 Write in brief (Any Five)

10

- (a) Lung abscess
- (b) Surgical anatomy of the lungs
- (c) Triangle of safety.
- (d) Infective endocarditis,
- (e) Pericardial effusion
- (f) Indications & contraindications of heart transplantation.

4 Select the most appropriate answers from the options given below each question:

(all questions are compulsory)

- 1. Commonest cause of spontaneous pneumothorax is
 - (a) Trauma
 - (b) Rupture of sub-pleural bleb
 - (c) Tuberculosis
 - (d) Bronchial asthma
- 2. Bronchial tree is visualized with endoscope is called
 - (a) Bronchoscopy
 - (b) Mediastinoscopy
 - (c) Laparoscopy
 - (d) Thoracoscopy
- 3. Major cause of morbidity and mortality in developed country is due to
 - (a) Infective Heart disease
 - (b) Congenital Heart disease
 - (c) Rheumatic Heart Disease
 - (d) Ischemic Heart Disease
- 4. Which rib fracture alert the clinician to a potentially serious injury ?
 - (a) First rib
 - (b) Fifth rib
 - (c) Fourth rib
 - (d) Seventh rib
- 5. Most common mediastinal tumor is
 - (a) Lymphoma
 - (b) Germ cell tumour
 - (c) Thymoma
 - (d) Mesenchymal tumour
- 6. An empyema of the underlying thoracic cavity may discharge through the chest wall leaving a chronic sinus, this condition is known as
 - (a) Empyema necessitans
 - (b) Fibrothorax
 - (c) Bronciatesis
 - (d) Pectus excavatum

		(d)	anterior interosseous ne	erve	palsy		
		(c)	median nerve palsy				
		(b)	radial nerve palsy				
			ulnar nerve palsy				
	1. Wrist drop is due to						
8	Sele	ct cor	rect options in following	,		10	
	6.	Rota	tor cuff tear of shoulder	•			
	5.		is elbow				
	4.	Ricke	ets				
	3.	Foot	drop				
	2.	Ulna	r nerve palsy				
	1.	Dequiervan's disease of thumb					
7	Writ	Write in short (five out of six) 10					
	3.	Scur	vy				
	2.	Rheu	matoid arthritis				
•	1.						
6	Writ	ite in short about following any two out of three. 10					
	3.	Descr	ribe in detail about rad	ial n	erve palsy.		
	2.	Descr	ribe in detail about lum	bar	disc prolapsed		
	1.	Descr					
5	Write long assay on any two out of three.					20	
			SECTION-2 ORTH	UPA	EDICS		
		` /	Open cardiac massage	` '	Q		
			Wait and watch	(b)			
	10.	` /	rgency treatment of per		- -		
		` /	Pulmonary Stenosis	` /	Ventricular hy	nartronhy	
	9.		ch anomaly is not includ VSD	aea 1 (b)	n Tetralogy of a	fallot' ?	
	0	` ′	Leiomyoma	(d)	Lipoma	C 11 O	
		` /	Rhabdomyoma	(b)	Myxoma		
	8.		Most common primary caridac tumour is				
		(c)	Chest drain insertion	(d)	Rib resection		
		(a)	Wait and watch.	(b)	Thoracoscopy		
	7.	Treat	Treatment of traumatic pneumothorax is				

2.	Foot drop is caused by							
	(a) common peroneal nerve injury							
	(b) median nerve injury							
	(c) L 2 root injury							
	(d) L 3 root injury							
3.	Shoulder abduction is lost in							
	(a)	C 1 root injury	(b)	C 2 root injury				
	(c)	C 4 root injury	(d)	C 5 root injury				
4.	Jersey finger is injury of							
	(a)	FCU tendon	(b)	FCR tendon				
	(c)	FDS tendon	(d)	FDP tendon				
5.	March fracture is common in							
	(a)	swimmers	(b)	computer operators				
	(c)	traffic police	(d)	soldiers				
6.	Movie sign is due to							
	(a)	hip arthritis	(b)	patellofemoral arthritis				
	(c)	ankle arthritis	(d)	subtalar arthritis				
7.	Aviator's astralgus is							
	(a)	femur fracture	(b)	acetabulum fracture				
	(c)	upper tibia fracture	(d)	talus fracture				
8.	Game keeper's thumb is due to							
	(a) FPL injury							
	(b) EPL injury							
	(c) UCL injury of MCP joint							
	(d) RCL injury of MCP joint							
9.	Commonest nerve injury in supra condylar fracture of							
	humerus is							
	(a)	ulnar nerve	(b)	radial nerve				
	(c)	median nerve	(d)	axillary nerve				
10.	Harrison's sulcus is seen in							
	(a)	gout	(b)	rickets				
	(c)	rheumatoid arthritis	(d)	Goucher's disease				