



PAF-003-010401 Seat No. _____

M. Sc. (Sem. IV) (CBCS) Examination

August – 2020

C - 401 : Spectroscopy

(Multidisciplinary) (Old Course)

Faculty Code : 003

Subject Code : 010401

Time : $2\frac{1}{2}$ Hours] [Total Marks : 70


Instruction : All questions are **compulsory** and carry **equal marks**.

- 1** Answer the following : (any seven) 14

 - (a) What is an IR Spectrum ? When does a molecule absorb IR light ?
 - (b) What is Nitrogen rule ? Validate this rule using nitrobenzene and 2,4-dinitrobenzene.
 - (c) Discuss types of UV absorption shift.
 - (d) Write the characteristic IR frequency of m-Chloro benzaldehyde.
 - (e) Distinguish between germinal and vicinal coupling with suitable example.
 - (f) Discuss the advantage of ^{13}C MR spectroscopy.
 - (g) Give the full-form of HMQC, HMBC, NOESY and TOCSY.
 - (h) Draw the ^1H NMR spectrum of p-Nitro toluene with splitting of the signals.
 - (i) Distinguish methyl propionate and ethyl acetate with the help of ^1H NMR spectroscopy.
 - (j) Write the principle of Mass-spectroscopy.

2 Answer the followings : (any two) 14

 - (a) Predict the ^{13}C NMR signal for the following :



- (b) Sketch the NMR spectrum of AMX and A₂B₂ system with suitable example.
- (c) Describe HSQC 2D NMR technique with suitable example.

3 Answer the following : 14

- (a) Write a note on Sample handling technique in IR spectroscopy.
- (b) Determine the molecular formula of a compound which gives the data in its mass spectrum.

m/z	120	130	131	132	133	134	135	136	137
Rel. abundance	30	100	31	98	12	32	1.7	3.5	0.07

OR

- (a) Draw the schematic diagram of Mass spectrophotometer and discuss it in detail.
- (b) Explain the Mass fragmentation of Methyl butyrate, Methyl benzoate and Acetophenone.

4 Answer the following : 14

- (a) Calculate ¹³CNMR chemical shifts for the following :
 - (i) HO—CH₂—CH₂—CH₂—CH₂—CH₂—NH₂
 - (ii) CH=C—CH₂—CH₂—CH₂—CH₃
 - (iii) 2-Ethylhexanol
- (b) Discuss McLafferty rearrangement with at least three different compounds.

5 Answer the following : (any two) 14

- (a) Discuss the (n+l) rule, its failure and solution with suitable example.
- (b) Give the application of IR-spectroscopy.
- (c) Draw the single beam UV-spectrophotometer and functioning it.
- (d) What are the empirical rules for calculation of absorption maxima in case of polyenes ?

Data-table for the calculation of ^{13}C Chemical shift

TABLE 5.1

^{13}C Atoms	Shift (ppm) (Δ)
α	+9.1
β	+9.4
γ	-2.5
δ	+0.3
ϵ	+0.1
1° (3°)*	-1.1
1° (4°)*	-3.4
2° (3°)*	-2.5
2° (4°)	-7.2
3° (2°)	-3.7
3° (3°)	-9.5
4° (1°)	-1.5
4° (2°)	-8.4

*The notations 1° (3°) and 1° (4°) denote a CH_3 group bound to a R_2CH group and to a R_3C group, respectively. The notation 2° (3°) denotes a CH_2 group bound to a R_2CH group, and so on.

TABLE 5.2

Compound	C-1	C-2	C-3	C-4	C-5
Methane	-2.3				
Ethane	5.7				
Propanes	15.8	16.3	15.8		
Butane	13.4	25.2	25.2		
Pentane	13.9	22.8	34.7	22.8	13.9
Hexane	14.1	23.1	32.2	32.2	23.1
Heptane	14.1	23.2	32.6	29.7	32.6
Octane	14.2	23.2	32.6	29.9	29.9
Nonane	14.2	23.3	32.6	30.0	30.3
Decane	14.2	23.2	32.6	31.1	30.5
Isobutane	24.5	25.4			
Isopentane	22.2	31.1	32.0	11.7	
Isohexane	22.7	28.0	42.0	20.9	14.3
Neopentane	31.7	26.1			
2,2-Dimethylbutane	29.1	30.6	36.9	8.9	
3-Methylpentane	11.5	29.5	36.9	(18.8, 3- CH_3)	
2,3-Dimethylbutane	19.5	34.3			
2,2,3-Trimethylbutane	27.4	33.1	38.3	16.1	
2,3-Dimethylpentane	7.0	25.3	36.3	(14.6, 3- CH_3)	

TABLE 5.3

Incremental Substituent Effects (ppm) on Replacement of H by Y In Alkanes. Y Is Terminal or Internal* (+ downfield, - upfield)

Y	α		β		γ	
	Terminal	Internal	Terminal	Internal	Terminal	Internal
CH_3	+ 9	+ 6	+ 10	+ 8	-2	
CH_2CH_3	+ 4.5		+ 6		-0.5	
COOH	+21	+16	+ 3	+ 2	-2	
COO^-	+25	+20	+ 5	+ 3	-2	
COOR	+20	+17	+ 3	+ 2	-2	
COCl	+33	+28			+ 2	
CONH_2	+22		+ 2.5		-0.5	
COR	+30	+24	+ 1	+ 1	-2	
CHO	+31		0		-2	
Phenyl	+23	+17	+ 9	+ 7	-2	
OH	+48	+41	+10	+ 8	-5	
OR	+58	+51	+ 8	+ 5	-4	
OCOR	+51	+45	+ 6	+ 5	-3	
NH_2	+29	+24	+11	+10	-5	
NH^+	+26	+24	+ 8	+ 6	-5	
NHR	+37	+31	+ 8	+ 6	-4	
NR_2	+42		+ 6		-3	
NR_3	+31		+ 5		-7	
NO_2	+63	+57	+ 4	+ 4	-3	
CN	+ 4	+ 1	+ 3	+ 3	-3	
SH	+11	+11	+12	+11	-4	
SR	+20		+ 7		-3	
F	+68	+63	+ 9	+ 6	-4	
Cl	+31	+32	+11	+10	-4	
Br	+20	+25	+11	+10	-3	
I	- 6	+ 4	+11	+12	-1	

*Add these increments to the shift values of the appropriate carbon atom in Table 5.2 or to the shift value calculated from Table 5.1.

Source: F.W. Wehrli, A.P. Marchand, and S. Wehrli, *Interpretation of Carbon-13 NMR Spectra*, 2nd ed., London: Heyden, 1983.

Table 3.14 δ values for the carbons in aromatic and heterocyclic molecules (sp^2 and sp^3 carbons listed)




TABLE 5.9

Substituent	C-1 (Attachment)	C-2	C-3	C-4	C of Substituent (ppm from TMS)
H	0.0	0.0	0.0	0.0	
CH ₃	9.3	+0.7	-0.1	-2.9	21.3
CH ₂ CH ₃	+15.6	-0.5	0.0	-2.6	29.2 (CH ₂), 15.8 (CH ₃)
CH(CH ₃) ₂	+20.1	-2.0	0.0	-2.5	34.4 (CH), 24.1 (CH ₃)
C(CH ₃) ₃	+22.2	-3.4	-0.4	-3.1	34.5 (C), 31.4 (CH ₃)
CH=CH ₂	+9.1	-2.4	+0.2	-0.5	137.1 (CH), 113.3 (CH ₂)
C≡CH	-5.8	+6.9	+0.1	+0.4	84.0 (C), 77.8 (CH)
C ₆ H ₅	+12.1	-1.8	-0.1	-1.6	
CH ₃ OH	+13.3	-0.8	-0.6	-0.4	64.5
CH ₂ OOCCH ₃	+7.7	-0.0	-0.0	-0.0	20.7 (CH ₂), 66.1 (CH ₃), 170.5 (C=O)
OH	+26.6	-12.7	+1.6	-7.3	
OCH ₃	+31.4	-14.4	+1.0	-7.7	54.1
OC ₆ H ₅	+29.0	-9.4	+1.6	-5.3	
OOCCH ₃	+22.4	-7.1	-0.4	-3.2	23.9 (CH ₃), 169.7 (C=O)
OCH	+8.2	+1.2	+0.6	+5.8	192.0
OOCCH ₃	+7.8	-0.4	-0.4	+2.8	24.6 (CH ₃), 195.7 (C=O)
OCC ₆ H ₅	+9.1	+1.5	-0.2	+3.8	196.4 (C=O)
OCCF ₃	-5.6	+1.8	+0.7	+6.7	
OCH	+2.9	+1.3	+0.4	+4.3	168.0
OOCCH ₃	+2.0	+1.2	-0.1	+4.8	51.0 (CH ₃), 166.8 (C=O) 168.5
OCCl	+4.6	+2.9	+0.6	+7.0	
C≡N	-16.0	+3.6	+0.6	+4.3	119.5
NH ₂	+19.2	-12.4	+1.3	-9.5	
N(CH ₃) ₂	+22.4	-15.7	+0.8	-11.8	40.3
NHCCH ₃	+11.1	-9.9	+0.2	-5.6	
NO ₂	+19.6	-5.3	+0.9	+6.0	
N=C=O	+5.7	-3.6	+1.2	-2.8	129.5
F	+35.1	-14.3	+0.9	-4.5	
Cl	+6.4	+0.2	+1.0	-2.0	
Br	-5.4	+3.4	+2.2	-1.0	
I	-32.2	+9.9	+2.6	-7.3	
CF ₃	+2.6	-3.1	+0.4	+3.4	
SH	+2.3	+0.6	+0.2	-3.3	
SCH ₃	+10.2	-1.8	+0.4	-3.6	15.9
SO ₂ NH ₂	+15.3	-2.9	+0.4	+3.3	
Si(CH ₃) ₃	+13.4	+4.4	-1.1	-1.1	

*See D. E. Ewing, *Org. Magn. Reson.*, 12, 499 (1979) for chemical shifts of 709 monosubstituted benzenes.

Table 3.17 δ values for the carbons in carbonyl groups and some other multiple-bonded environments (solvent shifts, ± 2 ppm, are commonly observed for $C=O$)

$R-CHO$	aliphatic aldehydes	200–205
$Ar-CHO$	aryl and conjugated aldehydes*	190–194
$\begin{array}{c} \diagup \\ Ar-CO-R' \\ \diagdown \end{array}$	dialkyl ketones	205–218
$Ar-CO-Ar$	aryl and conjugated ketones	196–199
$\begin{array}{c} \diagup \\ Ar-CO-Ar \\ \diagdown \\ CO \end{array}$		
	cyclohexanone derivatives	209–213
	cyclopentanone derivatives	214–220
	bicyclic ketones	215–219
	simple quinones	180–187
$R-COOH$	carboxylic acids and salts	166–181
$R-COO^+$		
$R-COO-R'$	aliphatic esters	169–176
$Ar-COOAr$	esters with conjugation in the acid or alcohol moiety	164–169
$\begin{array}{c} \diagup \\ Ar-COO \\ \diagdown \end{array}$		
	lactones	170–178
$-CO-O-CO-$	anhydrides, all classes	163–175
$-CON-$	amides, all classes including lactams	162–179

Table 3.17 (continued)

$-CO-NH-CO-$	imides	168–184
$-COCl$	acyl chlorides, all classes	167–172
$-NH-CO-NH-$	ureas	153–163
$-O-CO-O-$	carbonates	152–156
$R-C\equiv N$	nitriles	114–124
$R-N\equiv C$	isonitriles	156–158
$R-N=C=O$	isocyanates	120–130
$\begin{array}{c} \diagup \\ >C=NOH \\ \diagdown \end{array}$	oximes	148–158
$\begin{array}{c} \diagup \\ >C=NNHC CONH_2 \\ \diagdown \end{array}$	semicarbazones	158–160
$\begin{array}{c} \diagup \\ >C=N-NH \\ \diagdown \end{array}$	hydrazones	145–149
$\begin{array}{c} \diagup \\ >C=N-R \\ \diagdown \end{array}$	imines	157–175

*Note:

Table 3.15 Influence of functional group X on the chemical shift position (δ) of nearby carbons in alkane chains

X	$X-C-\underset{\alpha}{C}-\underset{\beta}{C}-\underset{\gamma}{C}$				
	α -shift			β -shift	γ -shift
	$X-CH_2-$	$X-CH-$ R	$X-C-$ R		
	1° or	2° or	3°		
$-CH_3$	9	6	3	9	-3
$-R$: see table 3.11					
{ axial $-CH_3$	1	-	-	5	-6
equatorial $-CH_3$	6	-	-	9	0
(in cyclohexanes)					
$-CH=CH_2$	22	16	12	7	-2
$-C\equiv CH$	4	-	-	3	-3
$-C_6H_5$, $-Ar$	23	17	11	10	-3
$-F$	70	-	-	8	-7
$-Cl$	31	35	42	10	5
$-Br$	19	28	37	11	-4
$-I$	-7 to 20	-	-	11	-2
$-NH_2$, $-NHR$, $-NR_2$	29	24	18	11	-4
$-NO_2$	62	-	-	3	-5
$-NHCOR$, $-NRCOR$	10	-	-	0	0
$-NH_3^+$	25	-	-	7	-3
$-CN$	3	4	-	2	-3
$-SH$	2	-	-	2	-2
$-OH$	50	45	40	9	-3
$-OR$	50	24	17	10	-6
$-OCOR$	52	50	45	7	-6
$-COOH$, $-COOR$, $-CON$	20	16	13	2	-3
$-COR$, $-CHO$	30	24	17	2	-3
$-SO_3H$, $-SO_2N$	50	--	--	3	0

Table 3.16 Influence of functional group X on the chemical shift positions (δ) of nearby carbons in alkene groups and benzene rings

Base values.	ethylene (δ 123)		and		benzene (δ 128)				
	C-1	C-2			X	C-1 (ipso)	ortho	meta	para
	Alkenes				Benzenes				
	C-1	C-2		(ipso)					
—CH ₃	10	-8	9			0	0	-2	
R,	16	-8	15			0	0	-2	
R,	23	-8	21			0	0	-2	
—CH=CH ₂	15	-6	9			0	0	-2	
—CH≡CH	—	—	-6			4	0	0	
—C ₆ H ₅ , —Ar	13	-11	13			-1	1	-1	
—F	25	-34	35			-14	1	-5	
—Cl	3	-6	6			0	1	-2	
—Br	-8	-1	-5			3	2	-2	
—I	-38	7	-32			10	3	-1	
—NH ₂	—	—	18			-13	1	-10	
—NHR	—	—	20			-14	1	-10	
—NR ₂	—	—	22			-16	1	-10	
—NO ₂	22	-1	20			-5	1	6	
—NHCOR, —NRCOR	—	—	10			-7	1	-4	
—CN	-15	15	-16			4	1	6	
—SH	—	—	4			1	1	-3	
—OH	—	—	27			-13	1	-7	
—OR	29	-39	30			-15	1	-8	
—OCOR	18	-27	23			-6	1	-2	
—COOH, —COOR, —CONH	4	9	2			2	0	5	
—COR, —CHO	14	13	9			1	1	6	
—SO ₃ H, —SO ₂ NH	—	—	16			0	0	4	
—PMe ₂	—	—	14			1.6	0	-1	
—PAR ₂	—	—	9			5	0	0	

Table 3.13 δ values for the carbons in alkynes (sp)

REPRESENTATIVE ALKYNES									
acetylene (ethyne) CH≡CH δ 72									
R—C≡CH	83	66	R—C≡C—R'	82	C ₆ H ₅ —C≡CH	83	78		
C ₆ H ₅ —C≡C—R	86				C ₆ H ₅ —C≡C—C ₆ H ₅	90			
EMPIRICAL PREDICTIONS FOR OTHER ALKYNE CARBONS									
Chemical shift (δ) = 72 + Σ (increments for carbon atoms)									
$\begin{array}{ccccccccc} \text{C} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{C} \\ & \delta & \gamma & \beta & \alpha & & \alpha' & \beta' & \gamma' \quad \delta' \\ \text{increments} & +0.5 & 0 & +5 & +7 & & -6 & +2 & -1 \quad +0.5 \\ & & & & & & \text{base value } 72 & & \end{array}$									